arXiv:gr-qc/0509011v1 2 Sep 2005

On Some Accelerating Cosmological Kaluza-Klein Models

Goran S. Djordjević and Ljubiša Nešić
Department of Physics, University of Nis, Serbia and Montenegro
E-mail: nesiclj, gorandj@junis.ni.ac.yu

Abstract. We consider (4+D)-dimensional Kaluza-Klein cosmological model with two scaling factors, as real, p-adic and adelic quantum mechanical one. One of the scaling factors corresponds to the D-dimensional internal space, and second one to the 4-dimensional universe. In standard quantum cosmology, i.e. over the field of real numbers R, it leads to dynamical compactification of additional dimensions and to the accelerating evolution of 4-dimensional universe. We construct corresponding p-adic quantum model and explore existence of its p-adic ground state. In addition, we explore evolution of this model and a possibility for its adelic generalization. It is necessary for the further investigation of space-time discreteness at very short distances, i.e. in a very early universe.

1 Introduction

On the basis of the results of quantum gravity [1], theoretical uncertainty of measuring distances is greater or equal to the Planck distance. It leads to the conclusion that on very short distances Archimedean axiom is not valid, i.e. the space can posses ultrametric features. Because geometry is always connected with a concrete number field, in the case of the nonarchimedean geometry it is used to be the field of p-adic numbers Q_p. In the high energy physics, these numbers have been used almost twenty years. The motivation comes from string theory [2]. Generally speaking, p-adic approach should be useful in describing a very early phase of the universe and processes around Planck scale.

A significant number of papers, motivated by [2], has been published up to now (for a review see [3, 4]). In this paper we are especially interested in application of p-adic numbers and analysis in quantum cosmology [5]. p-Adic quantum mechanics [6] (QM) has been successfully applied in minisuperspace quantum cosmology [7]. We have treated many cosmological models, mainly constructed in four space-time dimensions [5]. Based on that one can ask: is it possible to extend this approach to the multidimensional quantum cosmological models? In this article we briefly formulate such a model.

After brief mathematical introduction in Section 2, a short review of p-adic and adelic quantum mechanics and cosmology is given in Section 3. Section 4 is devoted to the classical (4+D)-dimensional cosmological models filed with "exotic" fluid [8]. The corresponding p-adic model is considered in Section 5. This paper is ended by short conclusion, including adelic generalization, and suggestion for advanced research.

2 p -Adic Numbers and Adeles

Perhaps the most easier way to understand p-adic numbers is if one starts with the notion of norm. It is well known that any norm must satisfy three conditions: nonnegativity, homogeneity and triangle inequality. The completion of the field of rational numbers Q with respect to the absolute value, or standard norm $|.|_\infty$, gives the field of real numbers $R \equiv Q_\infty$. Besides this norm there are another ones which satisfy the first two conditions and the third one in a stronger way

$$\|x + y\| \leq \max(\|x\|, \|y\|), \tag{1}$$

so called strong triangle inequality. The most important of them is p-adic norm $|.|_p$ [3]. The feature (1), also called ultrametricity, is one of the most important characteristics of the p-adic norm (p denotes a prime number). The number fields obtained by completion of Q with respect to this norm are called p-adic number fields Q_p. It is known that any nonzero rational number x can be expressed as $x = \pm p^\gamma a/b$, where a γ is the rational number, and a and b are the natural numbers which are not divisible with the prime number p and have no common divisor. Then, p-adic norm of x is, by definition, $|x|_p = p^{-\gamma}$.

Because Q_p is local compact commutative group the Haar measure can be introduced, which enables integration. In particular, the Gauss integral will be employed

$$\int_{Q_p} \chi_p(\alpha x^2 + \beta x)dx = \lambda_p(\alpha)|2\alpha|_p^{-1/2}\chi_p\left(-\frac{\beta^2}{4\alpha}\right), \quad \alpha \neq 0, \tag{2}$$

where $\chi_p = \exp 2\pi i\{x\}_p$, is an additive character ($\{x\}_p$ is the fractional part of x), and $\lambda_p(\alpha)$ is an arithmetic complex-valued function [3].

It should be noted that there is the real counterpart of λ_p

$$\lambda_\infty(\alpha) = \frac{1}{\sqrt{2}}(1 - i\mathrm{sign}\alpha), \quad \alpha \in Q_\infty. \tag{3}$$

Commonly the main properties of λ_v, (v denotes ∞ or any p) are:

$$\lambda_v(0) = 1, \quad \lambda_v(a^2\alpha) = \lambda_v(\alpha), \tag{4}$$

$$\lambda_v(\alpha)\lambda_v(\beta) = \lambda_v(\alpha + \beta)\lambda_v(\alpha^{-1} + \beta^{-1}), \quad |\lambda_v(\alpha)|_\infty = 1. \tag{5}$$

Very simple but rather important function in p-adic analysis and p-adic QM is

$$\Omega(|x|_p) = \begin{cases} 1, & |x|_p \leq 1, \\ 0, & |x|_p > 1, \end{cases} \tag{6}$$

which is the characteristic function of Z_p. Note that $Z_p = \{x \in Q_p : |x|_p \leq 1\}$ is the ring of p-adic integers.

Simultaneous treatment of real and p-adic numbers can be realized by concept of adeles. An adele $a \in \mathcal{A}$ is an infinite sequence $a = (a_\infty, a_2, \cdots, a_p, \cdots)$, where $a_\infty \in R$ and $a_p \in Q_p$, with the restriction that $a_p \in Z_p$ for all but a finite set S of primes p. The set of all adeles $\mathcal{A}$ can be written in the form

$$\mathcal{A} = \underset{S}{U}\mathcal{A}(S), \quad \mathcal{A}(S) = R \times \prod_{p \in S} Q_p \times \prod_{p \notin S} Z_p\ . \tag{7}$$

Also, $\mathcal{A}$ is a topological space. Algebraically, it is a ring with respect to the componentwise addition and multiplication. There is the natural generalization of analysis on R and Q_p to analysis on $\mathcal{A}$ [9].

3 p -Adic and Adelic Quantum Mechanics and Cosmology

p-Adic quantum mechanics is quantum mechanics over the field Q_p, and p-adic quantum cosmology is the application of p-adic QM on the universe as a whole [5].

3.1 p -Adic and Adelic Quantum Mechanics

p-Adic QM has been developed in two different ways: in the first one wave function is complex valued function of p-adic variable [3], and in the second one, p-adic wave function depends on p-adic variable [10]. If we want simultaneous (adelic) treating of standard quantum mechanics and all p-adic's, including usual probabilistic interpretation developed in standard QM, then the first formulation is preferable one.

Because p-adic fields Q_p are totally disconnected [3], dynamics of p-adic quantum models is described by a unitary evolution operator $U(t)$ without using the Hamilton operator and infinitesimal displacement. Therefore, $U(t)$ is formulated in terms of its kernel $\mathcal{K}_t(x,y)$

$$U_p(t)\Psi(x) = \int_{Q_p} \mathcal{K}_t(x,y)\Psi(y)dy. \tag{8}$$

In the above equation $\mathcal{K}_t$ is the kernel of the p-adic evolution operator which is defined by the functional integral (in units $G = \hbar = c = 1$)

$$\mathcal{K}_t(x,y) = \int \chi_p\left(-\int_0^t L(\dot{q},q)dt\right)\prod_t dq(t), \tag{9}$$

with properties analogous to the properties existing in standard QM. In that way, p-adic QM is given by a triple [3] $(L_2(Q_p), W_p(z_p), U_p(t_p))$. Keeping in mind that standard QM can be also presented as the corresponding triple, ordinary and p-adic QM can be unified in the form of adelic quantum mechanics [11]

$$(L_2(\mathcal{A}), W(z), U(t)), \tag{10}$$

where $L_2(\mathcal{A})$ is the Hilbert space on $\mathcal{A}$, $W(z)$ is the unitary representation of the Heisenberg-Weyl group on $L_2(\mathcal{A})$ and $U(t)$ is the unitary representation of the evolution operator on $L_2(\mathcal{A})$.

The adelic evolution operator $U(t)$ is defined as

$$U(t)\Psi(x) = \int_{\mathcal{A}} \mathcal{K}_t(x,y)\Psi(y)dy = \prod_v \int_{Q_v} \mathcal{K}_t^{(v)}(x_v,y_v)\Psi^{(v)}(y_v)dy_v. \tag{11}$$

The eigenvalue problem for $U(t)$ reads

$$U(t)\Psi_{\alpha\beta}(x) = \chi(E_\alpha t)\Psi_{\alpha\beta}(x), \tag{12}$$

where $\Psi_{\alpha\beta}$ are adelic eigenfunctions, $E_\alpha = (E_\infty, E_2, ..., E_p, ...)$ denotes adelic energy, indices α and β denote energy levels and their degeneration, respectively. Any adelic eigenfunction has the form

$$\Psi_S(x) = \Psi_\infty(x_\infty)\prod_{p\in S}\Psi_p(x_p)\prod_{p\notin S}\Omega(|x_p|_p), \quad x \in \mathcal{A}, \tag{13}$$

where $\Psi_\infty \in L_2(R)$ and $\Psi_p \in L_2(Q_p)$ are ordinary and p-adic eigenfunctions, respectively. We would like to paraphrase Manin's opinion [12] that our universe cannot be described fully by real numbers nor p-adic's only, than by adeles. In other words, our universe is inherently adelic.

3.2 p -Adic and Adelic Quantum Cosmology

Adelic quantum cosmology can be understood as an application of adelic quantum theory to the universe as a whole [5]. In the path integral approach to standard quantum cosmology, the starting point is Feynman's path integral method, i.e. the amplitude to go from one state with intrinsic metric h'_{ij} and matter configuration ϕ' on an initial hypersurface Σ' to another state with metric h''_{ij} and matter configuration ϕ'' on a final hypersurface Σ'' is given by a functional integral

$$\langle h''_{ij}, \phi'', \Sigma'' | h'_{ij}, \phi', \Sigma' \rangle_\infty = \int \mathcal{D}(g_{\mu\nu})_\infty \mathcal{D}(\Phi)_\infty \chi_\infty(-S_\infty[g_{\mu\nu}, \Phi]), \qquad (14)$$

over all (four)-geometries $g_{\mu\nu}$ and matter configurations Φ, which interpolate between the initial and final configurations [13]. In above expression $S[g_{\mu\nu}, \Phi]$ is an Einstein-Hilbert action for the gravitational and matter fields.

To perform p-adic and adelic generalization we first make p-adic counterpart of the action using form-invariance under the inter-change of the real to the p-adic number fields. Then we generalize (14) and introduce p-adic complex-valued cosmological amplitude

$$\langle h''_{ij}, \phi'', \Sigma'' | h'_{ij}, \phi', \Sigma' \rangle_p = \int \mathcal{D}(g_{\mu\nu})_p \mathcal{D}(\Phi)_p \chi_p(-S_p[g_{\mu\nu}, \Phi]), \qquad (15)$$

where $g_{\mu\nu}(x)$ and $\Phi(x)$ are the corresponding p-adic counterparts of metric and matter fields continually connecting their values on Σ' and Σ''.

The standard minisuperspace ground state wave function in the Hartle-Hawking (no-boundary) proposal [14], will be attained if one performs a functional integration in the Euclidean version of

$$\Psi_\infty[h_{ij}] = \int \mathcal{D}(g_{\mu\nu})_\infty \mathcal{D}(\Phi)_\infty \chi_\infty(-S_\infty[g_{\mu\nu}, \Phi]), \qquad (16)$$

over all compact four-geometries $g_{\mu\nu}$ which induce h_{ij} at the compact three-manifold. This three-manifold is the only boundary of all the four-manifolds. If we generalize the Hartle-Hawking proposal to the p-adic minisuperspace, then an adelic Hartle-Hawking wave function is an infinite product

$$\Psi[h_{ij}] = \prod_v \int \mathcal{D}(g_{\mu\nu})_v \mathcal{D}(\Phi)_v \chi_v(-S_v[g_{\mu\nu}, \Phi]), \qquad (17)$$

where the path integration must be performed over both, archimedean and nonarchimedean, geometries. If evaluation of the corresponding functional integrals gives as a result $\Psi[h_{ij}]$ in the form (13), we will say that such cosmological model is adelic one.

A more successful p-adic generalization of the minisuperspace cosmological models can be performed in the framework of p-adic and adelic quantum mechanics without using the Hartle-Hawking proposal. In such cases, we examine the conditions under which some eigenstates of the evolution operator exist [5].

3.3 Quantum Minisuperspace Models in p-Adic and Adelic Quantum Mechanics

In the study of the homogeneous universes, the metric depends only on the time parameter. It enables formulation of the models with a finite dimensional configuration space, so called minisuperspace [13]. Their variables are the three-metric components, and corresponding material fields, if any. The quantization of these models can be performed adopting of the rules of QM. One of the main reasons for using minisuperspace models in describing the universe is their simplicity. While the full (superspace) quantum cosmological models are usually unsolvable, the minisuperspace ones can be handled with available mathematical tools.

In p-adic minisuperspace quantum cosmology, we investigate necessary conditions for the existence of the p-adic ground state in the form of Ω-function. Some other typical eigenfunctions like δ-function can be also of interest. The existence of these functions enables construction of p-adic models and their adelization. Usually, this leads to the some restrictions on the parameters for the many exactly soluble minisuperspace cosmological models [5].

The necessary condition for the existence of an adelic quantum model is the existence of p-adic ground state $\Omega(|q_\alpha|_p)$ defined by (6), i.e.

$$\int_{|q_\alpha{}'|_p\leq 1} \mathcal{K}_p(q_\alpha{}'', N; q_\alpha{}', 0)dq_\alpha{}' = \Omega(|q_\alpha{}''|_p), \tag{18}$$

(q_α are minisuperspace coordinates, $\alpha = 1, 2, ..., n$ and N is the lapse function). Analogously, if a system is in the state $\Omega(p^\nu|q_\alpha|_p)$, where $\Omega(p^\nu|q_\alpha|_p) = 1$ if $|q_\alpha|_p \leq p^{-\nu}$ and $\Omega(p^\nu|q_\alpha|_p) = 0$ if $|q_\alpha|_p > p^{-\nu}$, then its kernel must satisfy equation

$$\int_{|q_\alpha{}'|_p\leq p^{-\nu}} \mathcal{K}_p(q_\alpha{}'', N; q_\alpha{}', 0)dq_\alpha{}' = \Omega(p^\nu|q''_\alpha|_p). \tag{19}$$

If p-adic ground state is of the form of the δ-function, where δ-function is defined as

$$\delta(a-b) = \begin{cases} 1, & if \quad a = b, \\ 0, & if \quad a \neq b, \end{cases}$$

then the corresponding kernel of the model has to satisfy equation

$$\int_{|q'_\alpha|_p = p^\nu} \mathcal{K}_p(q''_\alpha, T; q', 0)dq'_\alpha = \delta(p^\nu - |q''_\alpha|_p), \;\; \nu \in Z. \tag{20}$$

Equations (18) and (19) are usual p-adic vacuum ingredients of the adelic eigenvalue problem (12), where $\chi_p(Et) = 1$ in the vacuum state $\{Et\}_p = 0$. The above Ω and δ functions do not make a complete set of p-adic eigenfunctions, but they are very simple and illustrative. Since these functions have finite supports, the ranges of integration in (18)-(20) are also finite. The lapse function N is under the kernel $\mathcal{K}(q''_\alpha, N; q'_\alpha, 0)$ and N is restricted to the some values on which eigenfunctions do not depend explicitly.

4 (4+D)-Dimensional Cosmological Models Over the Field of Real Numbers

The old idea that four dimensional universe, in which we exist, is just our observation of physical multidimensional space-time attracts much attention nowadays. In such models compactification of extra dimensions play the key role and in the some of them leads to the period of accelerated expansion of the universe [8, 15, 16]. This approach is supported and encouraged with the recent results of the astronomical observations. We briefly recapitulate some facts of the real multidimensional cosmological models, necessary for p-adic and adelic generalization. The metric of such Kaluza-Klein model with D-dimensional internal space can be presented in the form [8, 17]

$$ds^2 = -\tilde{N}^2(t)dt^2 + R^2(t)\frac{dr^i dr^i}{(1+\frac{kr^2}{4})^2} + a^2(t)\frac{d\rho^a d\rho^a}{(1+k'\rho^2)^2} \tag{21}$$

where $\tilde{N}(t)$ is a lapse function, $R(t)$ and $a(t)$ are the scaling factors of 4-dimensional universe and internal space, respectively; $r^2 \equiv r^i r^i (i = 1, 2, 3)$, $\rho^2 \equiv \rho^a \rho^a (a = 1, ...D)$, and $k, k' = 0, \pm 1$. The form of the energy-momentum tensor is

$$T_{AB} = diag(-\rho, p, p, p, p_D, p_D, ..., p_D), \tag{22}$$

where indices A and B run over both spacetime coordinates and the internal space dimensions. If we want the matter is to be confined to the four-dimensional universe, we set all $p_D = 0$.

Now, we examine the case for which the pressure along all the extra dimensions vanishes $p_D = 0$ (in braneworld scenarios the matter is to be confined to the four-dimensional universe), so that all components of T_{AB} are set to zero except the spacetime components [8]. We assume the energy-momentum tensor of spacetime to be an exotic fluid χ with the equation of state

$$p_\chi = \left(\frac{m}{3} - 1\right)\rho_\chi, \tag{23}$$

(p_χ and ρ_χ are the pressure and the energy density of the fluid, parameter m has value between 0 and 2).

4.1 Classical Model

Dimensionally extended Einstein-Hilbert action (without a higher-dimensional cosmological term) is

$$S = \int \sqrt{-g}\tilde{R} dt d^3 R d^D \rho + S_m = \kappa \int dt L \tag{24}$$

where κ is an irrelevant constant, $\tilde{R}$ is the scalar curvature of the metric, so we can read off the lagrangian of the model (for flat internal space)

$$L = \frac{1}{2\tilde{N}} R a^D \dot{R}^2 + \frac{D(D-1)}{12\tilde{N}} R^3 a^{D-2} \dot{a}^2 + \frac{D}{2\tilde{N}} R^2 a^{D-1} \dot{R}\dot{a} - \frac{1}{2} k \tilde{N} R a^D + \frac{1}{6} \tilde{N} \rho_\chi R^3 a^D. \tag{25}$$

For closed universe ($k = 1$), substitution of the equation of state in the continuity equation

$$\dot{\rho}_\chi R + 3(p_\chi + \rho_\chi)\dot{R} = 0 \tag{26}$$

leads to the energy density in form

$$\rho_\chi(R) = \rho_\chi(R_0)\left(\frac{R_0}{R}\right)^m, \tag{27}$$

where R_0 is the value of scaling factor in arbitrary reference time. If we define cosmological constant as $\Lambda = \rho_\chi(R)$, lagrangian becomes

$$L = \frac{1}{2\tilde{N}} R a^D \dot{R}^2 + \frac{D(D-1)}{12\tilde{N}} R^3 a^{D-2} \dot{a}^2 + \frac{D}{2\tilde{N}} R^2 a^{D-1} \dot{R}\dot{a} - \frac{1}{2}\tilde{N} R a^D + \frac{1}{6}\tilde{N}\Lambda R^3 a^D. \tag{28}$$

Growth of the scaling factor R, according to (27) leads to the decrease of the cosmological constant by the relation

$$\Lambda(R) = \Lambda(R_0)\left(\frac{R_0}{R}\right)^m. \tag{29}$$

This decaying Λ term may also explain the smallness of the present value of the cosmological constant since, as the universe evolves form its small to large size, the large initial value of Λ decays to small values. If we take $m = 2$, initial condition for cosmological constant and scaling factor $\Lambda(R_0)R_0^2 = 3$, for laps function $\tilde{N}(t) = R^3(t)a^D(t)N(t)$, the Lagrangian (28) becomes

$$L = \frac{1}{2N}\frac{\dot{R}^2}{R^2} + \frac{D(D-1)}{12N}\frac{\dot{a}^2}{a^2} + \frac{D}{2N}\frac{\dot{R}\dot{a}}{Ra}. \tag{30}$$

It should be noted that there are no parameters k and Λ in the lagrangian. That means that although they are not zero in this model, it is equivalent to a flat universe with zero cosmological term. In another words, there is not difference between four dimensional universe which looks flat and not filled with fluid and closed universe filled with exotic fluid.

The solutions of corresponding equations of motion are

$$R(t) = C_1 e^{\alpha t}, \tag{31}$$

$$a(t) = C_2 e^{\beta t}, \tag{32}$$

where the constants C_1, C_2, α and β depend of initial conditions. It is a reasonable assumption that the size of all spatial dimensions is the same at $t = 0$. It may be assumed that this size would be the Planck size, i.e. $R(0) = a(0) = l_P$. Above solutions, can be read in terms of Huble parameter $H = \dot{R}/R$ [8]

$$R(t) = l_P e^{Ht}, \tag{33}$$

$$a(t) = l_P e^{-Ht}. \tag{34}$$

Depending of the dimensionality of the internal space we have,

$$R(t) = l_P e^{Ht}, \tag{35}$$

$$a_\pm(t) = l_P e^{\frac{2Ht}{D}[-1 \pm \sqrt{1 - \frac{2}{3}(1 - \frac{1}{D})}]^{-1}}, \tag{36}$$

for $D=1$ and

$$R_{\pm}(t) = l_P e^{\frac{D\beta t}{2}[-1\pm\sqrt{1-\frac{2}{3}(1-\frac{1}{D})}]}, \tag{37}$$

$$a(t) = l_P e^{\beta t}, \tag{38}$$

for $D>1$. The solution corresponding to $D=1$ predicts an accelerating (de Sitter) universe and a contracting internal space, with exactly the same rates. In the case $D>1$ analysis is complicated, but results are similar.

4.2 Quantum Model

Quantum solutions are obtained from the Wheeler-DeWitt equation

$$H\Psi(R,a) = 0, \tag{39}$$

where H is the Hamiltonian and Ψ is the wave function of the universe. For this model above equation is read in the new variables ($X=\ln R$ and $Y=\ln a$)

$$\left[(D-1)\frac{\partial^2}{\partial X^2} + \frac{6}{D}\frac{\partial^2}{\partial Y^2} - 6\frac{\partial}{\partial X}\frac{\partial}{\partial Y}\right]\Psi(X,Y) = 0. \tag{40}$$

By the new change $x = X\frac{3}{D+3} + Y\frac{D}{D+3}$, $y = \frac{X-Y}{D+3}$, Wheeler-DeWitt equation takes a simple form

$$\left[-3\frac{\partial^2}{\partial x^2} + \frac{D+2}{D}\frac{\partial^2}{\partial y^2}\right]\Psi(x,y) = 0. \tag{41}$$

Equation (41) has the four possible solutions [8]

$$\Psi_D^{\pm}(x,y) = A^{\pm}e^{\pm\sqrt{\frac{\gamma}{3}}x\pm\sqrt{\frac{\gamma D}{D+2}}y}, \tag{42}$$

$$\Psi_D^{\pm}(x,y) = B^{\pm}e^{\pm\sqrt{\frac{\gamma}{3}}x\mp\sqrt{\frac{\gamma D}{D+2}}y}, \tag{43}$$

where $A^{\pm}$ and $B^{\pm}$ are the normalization constants. It is possible to impose the boundary conditions to get a $\Psi_D(R,a)=0$. For the further details see [8].

5 (4+D)-Dimensional Model Over the Field of p-Adic Numbers

Consideration of (4+D)-dimensional Kaluza-Klein model over the field Q_p will be started from the lagrangian in the form (30). All quantities in this Lagrangian will be treated as the p-adic ones. Taking again replacement $X=\ln R$ and $Y=\ln a$, it becomes

$$L = \frac{1}{2N}\dot{X}^2 + \frac{D(D-1)}{12N}\dot{Y}^2 + \frac{D}{2N}\dot{X}\dot{Y}. \tag{44}$$

The corresponding p-adic equations of motion are

$$\ddot{X} + \frac{D}{2}\ddot{Y} = 0, \quad \ddot{X} + \frac{D-1}{3}\ddot{Y} = 0. \tag{45}$$

It is not difficult to see that above system can be rewritten in the following form

$$\ddot{X} = 0, \quad \ddot{Y} = 0. \tag{46}$$

If we omit from consideration pseudoconstant solutions and concentrate on the analytical ones we get $X(t) = C_1 t + C_2$, $Y(t) = C_3 t + C_4$. The calculation of the p-adic classical action gives

$$\bar{S}_p(X'', Y'', N; X', Y', 0)$$

$$= \frac{1}{2N}(X'' - X')^2 + \frac{D(D-1)}{12N}(Y'' - Y')^2 + \frac{D}{2N}(X'' - X')(Y'' - Y'). \tag{47}$$

Because this action is quadratic with respect to both variables X and Y, we can write down the kernel of p-adic operator of evolution [18]

$$\mathcal{K}_p(X'', Y'', N; X', Y', 0)$$

$$= \lambda_p\left(\frac{D(D+2)}{48N^2}\right)\left|\frac{D(D+2)}{12N^2}\right|_p \chi_p(-\bar{S}_p(X'', Y'', N; X', Y', 0)) \tag{48}$$

Let use again the change $x = X\frac{3}{D+3} + Y\frac{D}{D+3}$, $y = \frac{X-Y}{D+3}$ to separate variables and make further analyzes of this model rather simple. In these variables the classical action and the kernel of evolution operator read

$$\bar{S}_p(x'', y'', N; x', y', 0)$$

$$= \frac{1}{2N}(1 + \frac{D(D+5)}{6})(x'' - x')^2 - \frac{1}{2N}D(D+3)(y'' - y')^2, \tag{49}$$

$$\mathcal{K}_p(x'', y'', N; x', y', 0) = \lambda_p\left(\frac{6 + D(D+5)}{6}\right)\lambda_p\left(-\frac{D(D+3)}{12N}\right)$$

$$\times \left|\frac{D(D+3)}{2N^2}\left(1 + \frac{D(D+5)}{6}\right)\right|_p^{1/2} \chi_p(-\bar{S}_p(x'', y'', N; x', y', 0)). \tag{50}$$

Now, we can examine when p-adic wave function has the form corresponding to the simplest ground state [19]

$$\Psi_p(x, y) = \Omega(|x|_p)\Omega(|y|_p). \tag{51}$$

Putting kernel of the operator of evolution (50) in equation (18) we get that the required state exists if both conditions

$$|N|_p \leq \left|1 + \frac{D(D+5)}{6}\right|_p, \quad |N|_p \leq |D(D+3)|_p, \; p \neq 2, \tag{52}$$

are fulfilled. To answer the question: "can this conditions be useful in determination of dimensionality of the internal space?" needs further careful analysis.

Going back to the "old variables", p-adic ground state wave function for our model is

$$\Psi_p(x, y) = \Omega\left(\left|\left(1 - \frac{D}{D+3}\right)X + \frac{D}{D+3}Y\right|_p\right)\Omega\left(\left|\frac{X-Y}{D-3}\right|_p\right). \tag{53}$$

We can also write down the solutions in the variables R and a.

6 Conclusion

In this paper, we demonstrate how a p-adic version of the quantum (4+D)-Kaluza-Klein model with exotic fluid can be constructed. It is an exactly soluble model. From equations (42), (43) and (51), i.e. (53), it is possible to construct an adelic model too, i.e. model which unifies standard and all p-adic models [12]. The investigation of its possible physical implication and discreteness of space-time deserves much more attention and room.

Let us note that adelic states for the (4+D)-dimensional Kaluza-Klein cosmological model (for any D which satisfies (52)) exist in the form

$$\Psi_S(x,y) = \Psi^{\pm}_{D,\infty}(x_\infty, y_\infty) \prod_{p\in S} \Psi_p(x_p, y_p) \prod_{p\notin S} \Omega(|x_p|_p)\Omega(|y_p|_p), \tag{54}$$

where $\Psi^{\pm}_{D,\infty}(x_\infty, y_\infty)$ are the corresponding real counterparts of the wave functions of the universe. In the ground state wave functions $\Psi_p(x_p, y_p)$ are proportional to $\Omega(p^\nu|x_p|_p)\ \times\Omega(p^\mu|y_p|_p)$ or to $\delta(p^\nu - |x_p|)\delta(p^\mu - |y_p|)$. Adopting the usual probability interpretation of the wave function (54), we have

$$|\Psi_S(x,y)|^2_\infty = \left|\Psi^{\pm}_{D,\infty}(x_\infty, y_\infty)\right|^2_\infty \prod_{p\in S} |\Psi_p(x_p, y_p)|^2_\infty \prod_{p\notin S} \Omega(|x_p|_p)\Omega(|y_p|_p), \tag{55}$$

because $(\Omega(|x|_p))^2 = \Omega(|x|_p)$.

As a consequence of Ω-function properties, at the rational points x, y and in the (special) vacuum state ($S = \emptyset$, i.e. all $\Psi_p(x_p, y_p) = \Omega(|x_p|_p)\Omega(|y_p|_p)$, we find

$$|\Psi(x,y)|^2_\infty = \begin{cases} \left|\Psi^{\pm}_{D,\infty}(x,y)\right|^2_\infty, & x, y \in Z, \\ 0, & x, y \in Q\backslash Z. \end{cases} \tag{56}$$

This result leads to some discretization of minisuperspace coordinates x, y. Namely, probability to observe the universe corresponding to our minisuperspace model is nonzero only in the integer points of x and y. Keeping in mind that Ω-function is invariant with respect to the Fourier transform, this conclusion is also valid for the momentum space. Note that this kind of discreteness depends on adelic quantum state of the universe. When some p-adic states are different from $\Omega(|x|_p)\Omega(|y|_p)$ ($S \neq \emptyset$), then the above adelic discreteness becomes less transparent.

Performing the integration in (55) over all p-adic spaces, and having in mind that eigenfunctions should be normed to unity, one recovers the standard effective model over real space. However, if the region of integration is over only some parts of p-adic spaces, then the adelic approach manifestly exhibits p-adic quantum effects. Since the Planck length is here the natural one, the adelic minisuperspace models refer to the Planck scale.

Further investigations will be focused on determination of conditions for existence of the ground states in the form $\Omega(p^\nu|x|_p)\Omega(p^\mu|y|_p)$ and p-adic delta function. We should emphasize that investigation of dimensionality D of internal space from the conditions (52) and pseudoconstant solutions of the equation (46) deserves attention. It could additionally contribute to better understanding of the model, especially from its p-adic sector.

Acknowledgments

The authors would like to thank the organizers of the "2nd Conference on Theoretical Physics: Titeica-Markov Sympozium" for the kind invitation to give this talk. Lj. Nesic is thankful to D. Greku and M. Visinescu for their kind hospitality during the Conference. The research of both authors was supported by the Serbian Ministry of Science and Technology, Project No 1643.

References

[1] L. J. Garay, *Int. J. Mod. Phys.* **A10** (1995) 145.

[2] I. V. Volovich, *Theor. Math. Phys.* **71** (1987) 337.

[3] V. S. Vladimirov, I. V. Volovich and E. I. Zelenov, *p-Adic Analysis and Mathematical Physics*, World Scientific, Singapore, 1994.

[4] L. Brekke and P. G. O Freund, "p-Adic numbers in physics", *Phys. Rep.* **233** (1993) p. 1.

[5] G. S. Djordjevic, B. Dragovich, Lj. Nesic, I.V.Volovich, *Int. J. Mod. Phys.* **A 17** (2002) p. 1413, gr-qc/0105050.

[6] G. S. Djordjevic, B. Dragovich and Lj. Nesic, *Modern Physics Letters* **A 14** (1999) p. 317.

[7] B. Dragovich and Lj. Nesic, *Grav. Cosm.* **5** (1999) 222.

[8] F. Darabi, *Classical and Quantum Gravity*, **20** (2003) 3385.

[9] I. M. Gel'fand, M. I. Graev and I. I. Piatetskii-Shapiro, *Representation Theory and Automorphic Functions* Saunders, London, 1966.

[10] A. Khrennikov, *Non-Archimedean Analysis: Quantum Paradoxes, Dynamical Systems and Biological Models*, Kluwer Acad. Publ., Dordrecht, 1997.

[11] B. Dragovich, *Theor. Mat. Phys.* **101** (1994) 349.

[12] Yu. I. Manin, in *Conformal invariance and string theory*, Acad. Press, (1989) 293.

[13] D. L. Wiltshire, "An Introduction to Quantum Cosmology", gr-qc/0101003.

[14] J. Hartle and S. Hawking, *Phys. Rev.* **D28** (1983) 2960.

[15] P. K. Townsend, M. N. R. Wohlfarth, "Accelerating cosmologies from compactification", hep-th/0303097.

[16] S. Jalalzadeh, F. Ahmadi, H. R. Sepangi, "Multi-dimensional classical and quantum cosmology", hep-th/0308067.

[17] J. Wudka, *Phys. Rev.* **D35** (1987) 3255.

[18] G. S. Djordjevic, B. Dragovich, Lj. Nesic, Int. Conference: FILOMAT 2001, Nis, 26-29 August 2001. FILOMAT 15 (2001) 323, G. S. Djordjevic, B. Dragovich, *Mod. Phys. Lett.* A12 (1997) 1455.

[19] G. S. Dordjevic and Lj. D. Nesic, in *Proc. of the XI Congress of Physicists of Serbia and Montenegro*, Petrovac, SCG, 3-5. June 2004, 6/29.